

New earthworm species of the genus *Amyntas* Kinberg, 1867 from Thailand (Clitellata, Oligochaeta, Megascolecidae)

Ueangfa Bantaowong^{1,†}, Ratmanee Chanabun^{1,‡}, Piyoros Tongkerd^{1,§}, Chirasak
Sutcharit^{1,||}, Samuel W. James^{2,¶}, Somsak Panha^{1,#}

1 *Animal Systematics Research Unit, Department of Biology, Faculty of Science, Chulalongkorn University, 254 Phayathai Road, Pathumwan, Bangkok 10330, Thailand* **2** *Biodiversity Institute, University of Kansas, Lawrence, KS 66045, USA*

[†] urn:lsid:zoobank.org:author:11DD5A64-23CF-4618-A5EB-33454D590B03

[‡] urn:lsid:zoobank.org:author:024BBACF-15CE-4306-AF9D-CC01F9AFC8F9

[§] urn:lsid:zoobank.org:author:18575FA7-4812-4D75-8568-24D9E7D90BC6

^{||} urn:lsid:zoobank.org:author:AF740C51-9E43-4B72-904D-0DB5E8208791

[¶] urn:lsid:zoobank.org:author:7A2A4392-8B26-4E99-B26F-CE77AAFE81EF

[#] urn:lsid:zoobank.org:author:AC935098-D901-4F35-A414-4B0D4FE44E79

Corresponding author: Somsak Panha (somsak.pan@chula.ac.th)

Academic editor: Robert Blakemore | Received 11 February 2011 | Accepted 7 April 2011 | Published 14 April 2011

urn:lsid:zoobank.org:pub:81F631C6-D8FA-4301-B027-8C5CEBC21403

Citation: Bantaowong U, Chanabun R, Piyoros Tongkerd P, Sutcharit C, James SW, Panha S (2011) New earthworm species of the genus *Amyntas* Kinberg, 1867 from Thailand (Clitellata, Oligochaeta, Megascolecidae). ZooKeys 90: 35–62. doi: 10.3897/zookeys.90.1121

Abstract

Four new species of terrestrial earthworms from the *zebrus*-group in the genus *Amyntas* Kinberg, 1867, are described from Nan province, north Thailand: *A. phatubensis* **sp. n.**, from Tham Pha Tub Arboretum, *A. tontong* **sp. n.**, from Tontong Waterfall, *A. borealis* **sp. n.**, from Chaloeprakiat district, and *A. srinan* **sp. n.**, from Srinan National Park. After comparing with the two closely related Laos species *A. chandyi* Hong, 2008 and *A. namphouinensis* Hong, 2008, the four new species show clear morphological differences, and also it is confirmed that there are no previous records of the species described here. *Amyntas phatubensis* **sp. n.** is the largest (longest) sized of these earthworms and is the only species that lives in limestone habitats. The genital characters are different among them and also from the two Laotian species. Molecular systematics would be a good method for further analysis of the diversity and species boundaries in SE Asian *Amyntas*.

Keywords

Amyntas, Earthworm, Taxonomy, New species, Thailand

Introduction

Previous taxonomic publications on, or including, the Megascolecidae (*sensu* Blakemore 2000) of Thailand are comprised of those of Gates (1972), Sims and Easton (1972) and Blakemore (2006b, 2008, 2011) and Blakemore et al. (2007). Collectively, in these publications, 32 species are recorded for Thailand, belonging to five genera (*Amyntas* Kinberg, 1867, *Lampito* Kinberg, 1867, *Metaphire* Sims & Easton, 1972, *Polypheretima* Michaelsen, 1934 and *Perionyx* Perrier, 1872). The genus *Amyntas* is one of the dominant terrestrial earthworm genera that occurs throughout Thailand and nearby countries. From the classifications by Sims and Easton (1972) and reports by Blakemore (2006b, 2011) and Somniam (2008), it would seem that 14 species from this genus have been recorded from many areas in Thailand (Table 1). However, in addition Kosavitkul (2005) has reported six species of *Amyntas* from Khao Yai National Park, which included three unknown species, Chantaravisoot (2007) reported five species of *Amyntas* from various areas in Thailand that were all commented to be new to science, and Somniam (2008) recorded seven *Amyntas* species from Nakhonratchasima province of which many are still unidentified. Outside of Thailand, recent publications have included that by James (2004) who described a new species (*A. heaneyi*) from the Philippines; Shen and Yeo (2005) who reported four *Amyntas* species in Singapore, and Hong (2008) who described two species (*A. chandyi* and *A. namphouinensis*) from Laos, and also reported some publications by Vietnamese who studied the earthworm fauna in Laos and described *A. xuongmontis*. From the above data it is clear that there are still many species waiting to be discovered and described. The Animal Systematics Research Unit, Chulalongkorn University's members have surveyed terrestrial earthworms throughout Thailand since 2005 and a part of their results has been summarized in Chantaravisoot (2007). In the present paper we describe an additional four new species belonging to the *zebrus*-group, a provisional assemblage designated by Sims and Easton (1972). Each of these new species is known only from its type locality, but as more intensive collecting is undertaken in Thailand and other Asian countries, the known range and habitats of these species may be extended. The habitats of all four new species were in the topsoil layer covered with leaf litter of deciduous forests. The localities were in Nan province, in the north of Thailand, as shown in Figure 1.

Since none of the four species described in this paper seems to fit the descriptions of species described in the past, the purpose of this paper is to formally describe these species as new to science. Their descriptions follow.

Table 1. Morphological characteristics comparison of *Amyntas* species recorded in Thailand. The morphological characters are from the original description of each nominal species, except for the character with (*) are from Gates (1972). (**) indicate the known localities of *Amyntas* species in Thailand taken from Gates (1972), Kosavittikul (2005) and Somniyam (2008). Species group are as per Sims and Easton (1972)

Species	Species group	Body length (mm)	Number of segments	Spermathecal pores	Genital markings	Genital marking glands	Seminal vesicles	Prostate glands	Intestinal caeca	Distribution**
<i>A. hupbonensis</i> (Stephenson, 1931)	aeruginosus	225	142	7/8–8/9	absent	absent	large in XI, XII	XVI–XX	manicate, XXVII–	Chonburi
<i>A. alexandri</i> (Beddard, 1900)	corticis	145	133	5/6–8/9	absent	absent	XI, XII	XVII–XX	simple, XXVII–XX	Chiengrai, Chiangmai, Nakornratchasima, Bangkok, Chonburi
<i>A. comptus</i> (Gates, 1932)	corticis	197–260*	120–134*	5/6–8/9	three trios on 18/19–20/21	sessile	larger in XI, XII	XVIII	simple, XXVII–XXIII	Phrae
<i>A. exiguus austrinus</i> (Gates, 1932)	corticis	33–68	73–102	5/6–8/9	two pairs on 17/18, 18/19	absent	small in XI, XII	XVII–XX	simple, XXVII–XXIV	Chiangmai
<i>A. exiguus exiguus</i> (Gates, 1930)	corticis	43	90	5/6–8/9	paired on vii, viii, xix, xx	absent	small in XI, XII	XVII–XIX	simple, XXVII–XXIV	Phrae
<i>A. longicauliculatus</i> (Gates, 1931)	corticis	170	138	5/6–8/9	three pairs on 18/19–20/21	sessile	XI, XII	XVIII	simple, xxvii–xxiv	Chiangmai, Lumphun, Nakornratchasima
<i>A. manicata decorosa</i> (Gates, 1932)	corticis	40	60	5/6–8/9	one pair on xviii	sessile	large in XI, XII	XVII–XIX	manicate, XXVII–XXII	Chiangmai
<i>A. mekongianus</i> (Cognetti, 1922)	corticis	1 meter	370	5/6–8/9	absent	absent	10/11–11/12	XVII–XVIII	simple, XXVII–XXIII	Chiengrai

Species	Species group	Body length (mm)	Number of segments	Spermathecal pores	Genital markings	Genital marking glands	Seminal vesicles	Prostate glands	Intestinal caeca	Distribution**
<i>A. defecta</i> (Gates, 1930)	gracilis	>78	>49	5/6–7/8	absent	absent	small in XI, XII	absent	manicate, XXVII–XXXVI	Nakornratchasima
<i>A. gracilis</i> (Rosa, 1891)	gracilis	100	88–95	5/6–7/8	clusters on xviii	stalked*	XI, XII*	XVII–XXIII	simple, XXVII–XXIV*	Dor Kiu Koh Ma, north Thailand
<i>A. papulosus</i> (Rosa, 1896)	gracilis	45–50	110–115	5/6–7/8	transverse row on XVII–XIX	stalked*	XI, XII	XVI–XXI	simple, XXVII–XXII*	Yala
<i>A. morrissi</i> (Beddard, 1892)	morrissi	52	93	5/6–6/7	near spermathecal pore	stalked	XI, XII*	XVII–XXIII*	simple, XXVII–XXIV*	Chiangmai
<i>A. fucosus</i> (Gates, 1933)	sieboldi	120	114	6/7–8/9	two pairs on 17/18, 18/19	sessile	large in XI, XII	XVII–XX	simple, XXVII–XVII	Nakornratchasima
<i>Amyntas siam.</i> Blakemore, 2011	sieboldi	>70	>73	6/7–8/9	one pair postsetal on XVIII	sessile	XI, XII	XVIII–	simple, XXVII–	Sakon Nakhon




Figure 1. Map of type locality of **1** *Amyntas srinan* sp. n. from Srinan National Park, Nan province, **2** *Amyntas phatubensis* sp. n. from Tham Pha Tub Arboretum, Nan province, **3** *Amyntas tontong* sp. n. from Tontong Waterfall, Pua district, Nan province and **4** *Amyntas borealis* sp. n. from a small hill near Chaloemprakit district, Nan province.

Material and methods

Earthworms were collected from deciduous forests in many areas in Nan province, north of Thailand, by carefully digging up the topsoil near casts and by hand sorting the leaf litter. The worms were killed in 30% (v/v) ethanol, photographed, transferred to 5% (w/v) formalin for fixation for approximately 12 hours, and then transferred to 70% (v/v) ethanol for longer term preservation and subsequent morphological studies.

Duplicate specimens and/or tissue samples (in the cases of morphotypes determined to be unique on field inspection) were preserved in 95% ethanol for molecular data and DNA barcoding. Tissues were sent to the Canadian Center for DNA Barcoding (Hebert et al. 2003a, b) and processed according to their standard protocols (Hajibabaei et al. 2005; Ivanova et al. 2006; Ratnasingham and Hebert 2007). DNA barcode data are provided for paratype specimens of the first two species described in this paper. The sequences were aligned with Clustal X using default settings, and the resulting Neighbor-Joining tree (Saitou and Nei 1987) was used to identify barcode clusters. These clusters were matched to OTUs identified from quick examination of external characters. Inter- and intra- cluster genetic distances were calculated in MEGA

4 (Tamura et al. 2007) using the Kimura two parameter distance (Kimura 1980) using gamma-distributed rates among sites, pairwise deletion of sites with missing data, and using all substitution types and codon positions.

The descriptions of each species were made during observation under a Stemi DV 4 ZEISS stereoscopic light microscope. Drawings were made of the body segments and the distinct external characters and internal organs, as mentioned above, and are shown in Figures 2–5 for the four new species, respectively. The number of segments and the body width and length were measured in both full adults and juveniles, and are presented as the range (min-max) and mean±one standard deviation.

Type specimens housed at the Department of Biology, Faculty of Science, National University of Laos, Vientiane, Laos (BDNUL), of the two closely related Laos species, *A. chandyi* Hong, 2008 and *A. namphouinensis* Hong, 2008, have been critically studied and compared with the new species of this report.

Holotype and paratype specimens have been deposited in the Chulalongkorn University, Museum of Zoology, Bangkok, Thailand (CUMZ). Additional paratypes are housed in the Biozentrum Grindel und Zoologisches Museum, Hamburg, Germany (UHH), and the Natural History Museum, London (NHM).

Anatomical abbreviations: fp, female pore; ic, intestinal caeca; mp, male pores; pg, prostate gland; sc, spermathecae; sp, spermathecal pores; sv, seminal vesicles.

Systematics

Genus *Amyntas* Kinberg, 1867

Type species. *Amyntas aeruginosus* Kinberg, 1867, by monotypy.

Amyntas phatubensis Panha & Bantaowong, sp. n.

urn:lsid:zoobank.org:act:299379EB-C7CE-4B89-8A40-40E3122DCAB9

http://species-id.net/wiki/Amyntas_phatubensis

Figs 1, 2

Description of holotype: Dimensions; 110 mm by 4.3 mm at segment X, 4.3 at segment XX, 4.0 mm at clitellum; body cylindrical with 108 segments. Setae regularly distributed around segmental equators, numbering 51 at VII, 60 at XX, 15 between mp, setae formula AA:AB:ZZ:ZY= 1:1:1:1 at XIII with no ventral gaps. Single fp at XIV. Prostomium epilobic with tongue open. First dorsal pore at 5/6. Clitellum annular XIV–XVI with no setae.

A pair of mp is located ventro-laterally in XVIII, or at 9th seta line, 0.33 circumference apart ventrally, convex structure; distance between mp 4.2 mm. Porophores (protuberances bearing male aperture), papilla-like structures. Each mp surrounded by six flat, circular




Figure 2. External and internal morphology of holotype (CUMZ 3204) of *Amyntas phatubensis* sp. n. **A** External ventral view, **B** internal dorsal view and **C** spermatheca, and black arrow indicates the connection of the spermatheca and spermathecal pore.

genital markings almost the same diameter as mp, also one pair is equatorial in XVII in line with the male pores. One pair of sp in intersegmental furrow 7/8, distance between pores 0.32 circumference ventrally apart; distance between sp 3.5 mm. Genital markings, rounded, flat, located close to sp, postsetal paired on VII very near 7/8, presetal paired on VIII.

Septa 5/6 and 6/7 thick, 7/8 thin, 8/9 and 9/10 absent, 10/11–13/14 thin. Gizzard large within VIII–X, intestinal origin in XV, no lymph glands observed. Typhlosole small from XXVII. Intestinal caeca originate from XXVII extending forward to XXIII, simple, long finger-shape. Hearts esophageal in X–XIII. Holandric; testes and funnels in ventrally joined sacs in X–XI. Seminal vesicles paired in XI–XII. Prostates in XVII–XX; prostatic ducts U-shape. Genital marking glands absent.

Ovaries in XIII. Sc one pair in VIII; ampulla large ovate sac, duct stout, short; long stalked diverticulum, convoluted kinks enclosed within membrane, spherical knob terminal. No nephridia on spermathecal ducts. A large sessile genital marking gland corresponding to each external genital marking in VII–VIII.

All the key morphological characters of the holotype and paratype specimens are given in Table 2.

Variation: The holotype measures 110 mm body length with 108 segments; the twenty one paratypes range in size from 80–148 mm (108 ± 21.93 mm) body length with 85–114 segments (Table 2).

Type locality: Tham Pha Tub Arboretum, Nan province, Thailand, $18^\circ 51' 16.4''$ N, $100^\circ 44' 10.1''$ E, 265 meters elevation (11th October 2009). We also collected another lot of further specimens from Tontong Waterfall, Nan province (location 3 in Figure 1), which is located about a hundred kilometers north of the type locality.

Etymology: This species was named after the type locality, Tham Pha Tub Arboretum.

Type material: The holotype (CUMZ 3204) and 15 paratypes (CUMZ 3205) and 10 paratypes (CUMZ 3212) are deposited in Chulalongkorn University, Museum of Zoology. Another four paratypes will be deposited in the Biozentrum Grindel und Zoologisches Museum, Hamburg, Germany (UHH), and three paratypes in the Natural History Museum, London (NHM).

Habitat: Found in the top soil at about 10 cm depth, the soil surface was covered with leaf litter in a deciduous limestone forest at Tham Pha Tub Arboretum. The soil was carefully dug close to the casts. Many ariophantid snails, *Cryptozona siamensis* Pfeiffer, 1856 were on the ground or under leaf litter.

Diagnosis: *Amyntas phatubensis* sp. n. is a medium to large sized terrestrial earthworm with a pair of mp surrounded by six genital papillae on segment XVIII. Within the *zebrus*-group, this species is diagnosed by the unique combination of dorsal pores in 5/6, simple digitate caeca, ventrally joined testis sacs, genital marking glands in the spermathecal segments, and the spermathecal characters of the large ovate ampulla, stalked diverticulum whose folds are membrane-bound, and spherical knob terminal diverticulum sac.

Remarks: *Amyntas phatubensis* sp. n. has very simple characteristics of the genus, but among these, only the superficial male pores are external. In most newly collected specimens it was difficult to observe the pores or marks on the bodies. However, after preservation they can be seen more clearly. The internal organs are much more easily discerned. This new species is quite distinct when compared to the two closely related species from Laos, *A. chandyi* Hong, 2008 and *A. namphouinensis* Hong, 2008, which belong in the same *zebrus*-group. The two Laos species are a little bit smaller than *A. phatubensis* sp. n., especially *A. chandyi*. Even though *A. namphouinensis* is much closer in appearance to *A. phatubensis* sp. n., there are distinct differences between the type specimens (Figs 6 and 7). For example, the distance between the mp of *A. phatubensis* sp. n. is 4.2 mm for the holotype and range from 3.0–4.5 mm (4.27 ± 0.57 mm), while for *A. namphouinensis* this was significantly smaller, ranging from 1.4–1.5 mm. The distance between a pair of sp is also different, being 3.5–4.5 mm (4.12 ± 0.4 mm) for *A. phatubensis* sp. n. and 1.4–2.0 mm in *A. namphouinensis*. The distance between the male pores as a fraction of the estimated circumference of the 18th segment is 0.30–0.33 in *A. phatubensis* sp. n., but 0.10–0.14 circumference apart in *A. namphouinensis*.

Table 2. Holotype and Paratype dimension and other morphological characteristics of *Amyntas phatubensis* Panha & Bantaowong, sp. n.

Types	Characters	Body length (mm)	Number of segments	Location of genital markings		First dorsal pore	Number of setae			Prostate glands	Intestinal caeca
				Preclitellum	Postclitellum		VII	XX	Between male pore		
Holotype CUMZ 3204		110	108	VII, VIII	XVII, XVIII	5/6	51	60	15	XVII-XX	XXVII-XXIII
Paratype CUMZ 3205											
1		90	96	VII, VIII	XVIII	5/6	60	58	15	XVII-XXI	XXVII-XXIV
2		105	107	VII, VIII, IX	XVII, XVIII, XIX	5/6	52	58	12	XVII-XX	XXVII-XXIV
3		100	105	VII, VIII, IX	XVIII	5/6	53	60	9	XVII-XX	XXVII-XXIV
4		80	86	VII, VIII	XVII, XVIII, XIX	5/6	53	65	13	XVII-XX	XXVII-XXIV
5		120	96	VII, VIII	XVIII	5/6	58	68	11	XVII-XX	XXVII-XXIV
6		101	85	VII, VIII	XVIII	5/6	51	59	9	XVII-XX	XXVII-XXIV
7		131	86	VII, VIII	XVII, XVIII	5/6	64	67	15	XVII-XXI	XXVII-XXII
8		108	98	VII, VIII	XVII, XVIII	5/6	58	62	15	XVII-XXI	XXVII-XXII
9		116	99	VII, VIII	XVII, XVIII	5/6	53	64	11	XVII-XXI	XXVII-XXIII
10		89	92	VII, VIII	XVII, XVIII	5/6	64	58	12	XVII-XX	XXVII-XXIV
11		99	106	VII, VIII, IX	XVII, XVIII	5/6	60	63	13	XVII-XXI	XXVII-XXIV
12		112	112	VII, VIII	XVII, XVIII	5/6	52	58	11	XVII-XX	XXVII-XXIII
13		142	110	VII, VIII	XVII, XVIII	5/6	49	58	7	XVII-XX	XXVII-XXIV
14		137	108	VII, VIII, IX	XVII, XVIII	5/6	62	65	11	XVII-XX	XXVII-XXIII
15		80	85	VII, VIII, IX	XVII, XVIII	5/6	54	60	13	XVII-XX	XXVII-XXIV
16		89	111	VII, VIII, IX	XVII, XVIII	5/6	57	59	14	XVII-XXI	XXVII-XXIII
17		84	105	VII, VIII	XVII, XVIII	5/6	52	59	11	XVII-XX	XXVII-XXIV
18		148	112	VII, VIII	XVII, XVIII	5/6	51	58	12	XVII-XX	XXVII-XXII
19		109	114	VII, VIII	XVII, XVIII	5/6	64	59	12	XVII-XX	XXVII-XXII
20		144	107	VII, VIII	XVII, XVIII	5/6	53	60	11	XVII-XXI	XXVII-XXIV
21		84	108	VII, VIII	XVII, XVIII	5/6	64	61	15	XVII-XX	XXVII-XXII

Moreover, *A. phatubensis* sp. n. has no genital marking glands on segments XVII–XIX, where *A. namphouinensis* has sessile genital marking glands, but contains two distinct genital marking glands located close to sc that are absent in *A. namphouinensis*.

Two populations of *A. phatubensis* sp. n. were sampled, one from the type locality and one from Tontong waterfall. Distinct DNA barcode clusters corresponding to these populations had intra-cluster Kimura 2 parameter distances of 0.023 (N=9) and 0.016 (N=5) respectively. The inter-cluster divergence between the two populations is 0.084. Based on the morphological unity and the fact that the divergence is less than that usually seen between congeneric species pairs of earthworms (Chang et al. 2007; Pérez-Losada et al. 2005, James et al. 2010), we choose to maintain the two populations as representing one species. By contrast, the inter-cluster divergence between these populations and three other morpho-species with the same spermathecal battery, from the same two sites is in the range of 0.269–0.294. A consensus sequence from the type locality specimens is in Appendix 1. Another use of COI barcode sequence from type material is in Blakemore et al. (2010).

***Amyntas tontong* Panha & Bantaowong, sp. n.**

urn:lsid:zoobank.org:act:3317146B-143D-4EFC-A0C9-60262073BAFF

http://species-id.net/wiki/Amyntas_tontong

Figs 1, 3

Description of Holotype: Dimensions; 53 mm by 2.7 mm at segment X, 2.6 at segment XX, 2.2 mm at clitellum; body cylindrical with 80 segments. Setae regularly distributed around segmental equators, numbering 42 at VII, 52 at XX, no visible setae between mp, setae formula AA:AB:ZZ:ZY= 1.5:1:1:1 at XIII. Single fp at XIV. Prostomium epilobic. First dorsal pore at 5/6. Clitellum annular XIV–XVI with no setae.

A pair of indistinct rounded mp in XVIII, 0.19 mm circumference apart ventrally; distance between mp 1.0 mm at 5th seta line. Genital markings closely paired located medial to male pore level in intersegment 18/19. Sp paired in 7/8 at 4th seta line, each small, lip-like structure within porophore, 0.10 circumference apart ventrally; distance between sp 1.0 mm. Genital markings near sp absent.

Septa 5/6 and 6/7 thick, 7/8 thin, 8/9 and 9/10 absent 10/11–13/14 thin. Gizzard large within VIII–X, intestinal origin in XV, no lymph glands observed. Typhlosole small from XXVII. Ic originated from XXVII extending forward to XXV, simple finger-shape. Hearts esophageal in X–XIII. Holandric; testes and funnels in ventrally joined sacs in X–XI. Sv paired in XI–XII. Prostates in XVIII; prostatic ducts long slender with U-shape. Genital marking glands absent.

Ovaries in XIII. Sc one pair in VIII; ampulla thumb shape, duct stout, shorter than ampulla. Diverticulum slender stalk with spherical knob terminal, no genital marking glands observed.

All the key morphological characters of the holotype and paratype specimens are given in Table 3.




Figure 3. External and internal morphology of holotype (CUMZ 3206) of *Amyntas tontong* sp. n. **A** External ventral view, **B** internal dorsal view and **C** spermatheca, and black arrow indicates the connection of the spermatheca and spermathecal pore.

Variation: The holotype measures 53 mm body length with 80 segments; the three paratypes range in size from 39–41 mm (40.33 ± 1.15 mm) body length with 71–74 segments (Table 3).

Type locality: Tontong Waterfall, Nan province, Thailand, $19^{\circ}12'35.9''\text{N}$, $101^{\circ}04'13.7''\text{E}$, 1,128 meters elevation (10th October 2009).

Etymology: This species was named after the type locality, Tontong Waterfall.

Type material: The holotype (CUMZ 3206) and two paratypes (CUMZ 3207) are deposited in Chulalongkorn University, Museum of Zoology. Another paratype will be deposited in the Biozentrum Grindel und Zoologisches Museum, Hamburg, Germany (UHH).

Habitat: Found in the top soil at about 10 cm depth, the soil surface covered with leaf litter of deciduous forest which originated at the Tontong Waterfall area. The soil was carefully dug close to surface casts. Most surrounding areas have been modified to agricultural fields.

Table 3. Holotype and Paratype dimension and other morphological characteristics of *Amyntas tontong* Panha & Bantaowong sp. n.

Characters Types	Body length (mm)	Number of segments	Genital markings	First dorsal pore	Number of setae		Between male pore	Prostate glands	Intestinal caeca
					VII	XX			
Holotype CUMZ 3206	53	80	XVIII	5/6	42	52	0	XVII–XX	XXVII– XXIV
Paratype CUMZ 3207									
1	41	71	XVIII	5/6	41	53	0	XVI–XVIII	XXVII–XXV
2	39	74	XVIII	5/6	42	52	0	XVII–XX	XXVII– XXIV
3	41	73	XVIII	5/6	46	55	0	XVII–XIX	XXVII– XXIII

Diagnosis: *Amyntas tontong* sp. n. is a small sized terrestrial earthworm with a close indistinct pair of male pores with a pair of genital markings in intersegment 18/19. Spermathecae consists of a thumb shaped ampulla and a spherical terminal knob shaped diverticulum. Genital marking glands absent, first dorsal pore in 5/6, intestinal caeca simple, intestinal origin XV, septa 8/9/10 absent, testis sacs joined ventrally.

Remarks: *Amyntas tontong* sp. n., along with *A. srinan* sp. n. and *A. exiguus exiguus*, is one of the smallest sized *Amyntas* ever recorded in Thailand. The basic external characters are easily seen in both newly collected and preserved materials. Compared with the two other closely related species from Laos, *A. chandyi* Hong, 2008 and *A. namphouinensis* Hong, 2008, which belong in the same *zebrus*-group, *A. chandyi* is similar to *A. tontong* sp. n. However, it differs in the specific details of the significant characters, such as the distance between the mp in *A. tontong* sp. n. is 1.0 mm for the holotype and ranged from 1.0–1.2 mm (0.93 ± 0.12 mm), while in *A. chandyi* it ranged from 1.5–2.4 mm. The distance between the male pores as a fraction of the estimated circumference of the 18th segment is 0.15–0.19 in *A. tontong* sp. n., but 0.14–0.32 in *A. chandyi*. The arrangement of the genital markings of both species are totally different, and the distance between a pair of sp is also different, being 0.8–1.0 mm (1.1 ± 0.1 mm) in *A. tontong* sp. n. and 1.2–1.5 mm for *A. chandyi*. Moreover, *A. tontong* sp. n. has no genital markings near to the sp, whilst *A. chandyi* exhibits circular genital markings in various locations, paired or single mid ventral in VII, VIII; usually 3 or 4 in total.

Alcohol-preserved paratype specimens of *A. tontong* sp. n. belonged to a single DNA barcode cluster, with an intra-cluster divergence of 0.005 (N=3), and diverging from *A. phatubensis* sp. n. by 0.294, and by 0.189 for an undescribed species. An undescribed morph at Tham Pha Tub diverged by 0.100, and may represent a subspecies. A consensus sequence is in Appendix 1.

***Amyntas borealis* Panha & Bantaowong, sp. n.**

urn:lsid:zoobank.org:act:C2BE17F8-A721-4736-9809-EF9ABDAB0C03

http://species-id.net/wiki/Amyntas_borealis

Figs 1, 4

Description of Holotype: Dimensions; 54 mm by 3.5 mm at segment X, 3.8 at segment XX, 3.5 mm at clitellum; body cylindrical with 89 segments. Setae regularly distributed around segmental equators, numbering 39 at VII, 51 at XX, no visible setae between mp, setae formula AA:AB:ZZ:ZY= 2:1:1.5:1 at XIII. Single fp at XIV. Prostomium epilobic. First dorsal pore at 5/6. Clitellum annular XIV–XVI with no setae.

Mp pocket-like structures indistinctly occur in XVIII, 0.10 circumference apart ventrally; distance between mp 1.0 mm; porophores small, lip-like and surrounded by an elevated skin fold at medial pores, and there is a long ridge with a sharp posterior boundary traversing the body in front of the mp. Genital markings absent. Sp paired in 7/8 at 4th seta line, 0.10 circumference apart ventral; distance between sp 1.0 mm. Genital markings absent.

Septa 5/6 and 6/7 thick, 7/8 thin, 8/9 and 9/10 absent, 10/11–13/14 thin. Gizzard large within VIII–X, intestinal origin in XV, no lymph glands observed. Typhlosole small from XXVII. Ic originated from XXVII extending forward to XXV, simple finger-shape. Hearts esophageal in X–XIII. Holandric; testes and funnels in ventrally joined sacs in X–XI. Sv paired in XI–XII. Prostates in XVIII; prostatic ducts long slender bent in U-shape. Genital marking glands absent.

Ovaries at XIII. Sc one pair in VIII; ampulla large sac-shape, flattened by gizzard, narrow duct shorter than ampulla. Diverticulum with elongated tubular shape, stalk attached to duct near body wall, with no genital marking glands.

All the key morphological characters of the holotype and paratype specimens are given in Table 4.

Variation: The holotype measures 54 mm body length with 89 segments; the eight paratypes range in size from 42–45 mm (42.87 ± 1.25 mm) body length with 77–87 segments (Table 4).

Type locality: Chaloemprakiat district, Nan province, Thailand, 19°34'48.5"N, 101°04'53.1"E, 513 meters elevation (7th August 2010).

Etymology: The specific epithet “*borealis*” derived from Latin word “boreal” mean “north”. This name refers to the location of type locality in the north of Thailand.

Type material: The holotype (CUMZ 3208) and seven paratypes (CUMZ 3209) are deposited in Chulalongkorn University, Museum of Zoology. Another two paratypes will be deposited in the Biozentrum Grindel und Zoologisches Museum, Hamburg, Germany (UHH), and another two paratypes in the Natural History Museum, London (NHM).

Habitat: Found in the top soil at about 10 cm depth, the soil surface covered with the leaf litter of a deciduous limestone forest, mostly disturbed. The soil was carefully dug close to the casts.

Diagnosis: *Amyntas borealis* sp. n. is a small sized terrestrial earthworm small male pores, a transverse ridge anterior to the male pores in XVII, and no genital mark-




Figure 4. External and internal morphology of holotype (CUMZ 3208) of *Amynthes borealis* sp. n. **A** External ventral view, **B** internal dorsal view and **C** spermatheca, and black arrow indicates the connection of the spermatheca and spermathecal pore.

ings. One pair of sc in VIII, each spermathecae consists of a large sac-shaped ampulla and elongated tubular shaped diverticulum. Testis sacs joined ventrally, intestinal origin XV, intestinal caeca simple, first dorsal pore in 5/6.

Remarks: *Amynthes borealis* sp. n. is one of the smaller *Amynthes*. The characteristic male field is difficult to see in newly collected specimens but can be clearly observed after preservation. Compared with the two other closely related species from Laos, *A. chandyi* and *A. namphouinensis*, which belong in the same *zebrus*-group, *A. chandyi* is similar to *A. borealis* sp. n. However, distinctive differences include the distance between mp of the new species, being 1.0 mm in the holotype with a range of 0.8–1.0 mm (0.95 ± 0.09 mm) in *A. borealis* sp. n. compared to 1.5–2.4 mm. The distance between the male pores as a fraction of the estimated circumference of the 18th segment is 0.10–0.14 in *A. borealis* sp. n., but 0.14–0.32 in *A. chandyi*. There are no genital markings in the new species; the distance between a pair of sp is also different, being 0.5–1.0 mm (0.9 ± 0.19 mm) in the new species compared to 1.2–1.5 mm for *A. chan-*

Table 4. Holotype and Paratype dimension and other morphological characteristics of *Amyntas borealis* Panha & Bantaowong, sp. n.

Characters Types	Body length (mm)	Number of segments	Genital markings	First dorsal pore	Number of setae		Between male pore	Prostate glands	Intestinal caeca
					VII	XX			
Holotype CUMZ 3208	54	89	Absent	5/6	39	51	0	XVII–XIX	XXVII– XXV
Paratype CUMZ 3209									
1	45	87	Absent	5/6	51	48	0	XVII–XX	XXVII– XXIV
2	42	78	Absent	5/6	49	45	0	XVIII–XIX	XXVII– XXIII
3	44	79	Absent	5/6	51	50	0	XVII–XX	XXVII– XXIII
4	42	86	Absent	5/6	54	41	0	XVIII–XIX	XXVII– XXIV
5	44	85	Absent	5/6	40	40	0	XVIII–XIX	XXVII– XXIV
6	42	85	Absent	5/6	46	48	0	XVII–XIX	XXVII– XXIV
7	42	77	Absent	5/6	44	50	0	XVII–XX	XXVII– XXV
8	42	83	Absent	5/6	48	52	0	XVII–XIX	XXVII– XXV

dyi. Moreover, *A. borealis* sp. n. has no genital marking glands at all, whilst *A. chandyi* exhibits circular genital markings in various locations, paired or single mid ventral in VII and VIII; usually 3 or 4 in total.

***Amyntas srinan* Panha & Bantaowong, sp. n.**

urn:lsid:zoobank.org:act:C3EC91E6-B29A-4C72-908F-1858DE7F21DA

http://species-id.net/wiki/Amyntas_srinan

Figs 1, 5

Description of Holotype: Dimensions; 47 mm by 1.8 mm at segment X, 2.3 at segment XX, 2.3 mm at clitellum; body cylindrical with 77 segments. Setae regularly distributed around segmental equators, numbering 36 at VII, 42 at XX, four between mp, setae formula AA:AB:ZZ:ZY= 1.5:1:2:1 at XIII. Single fp at XIV. Prostomium epilobic with tongue open. First dorsal pore at 4/5 or 5/6. Clitellum annular XIV–XVI with no setae.

Mp on circular porophores in XVIII, 0.30 circumference apart ventrally; distance between mp 1.5 mm. Genital markings small, postsetal, closely paired near mid ventral




Figure 5. External and internal morphology of holotype (CUMZ 3210) of *Amynthes srinan* sp. n. **A** External ventral view, **B** internal dorsal view and **C** spermatheca, and black arrow indicates the connection of the spermatheca and spermathecal pore.

of XVII and XVIII. Sp paired in 7/8 at 6th setal lines, 0.26 circumference apart ventrally; distance between sp 1.5 mm. Genital markings tiny, closely paired on near mid ventral of VII and VIII.

Septa 5/6 and 6/7 thick, 7/8 thin, 8/9 and 9/10 absent, 10/11–13/14 thin. Gizzard globular within VIII–X, intestinal origin in XV, no lymph glands observed. Typhlosole small from XXVII. Ic originated from XXVII extending forward to XXIII, long finger-shape. Hearts esophageal in X–XIII. Holandric; testes and funnels in ventrally joined sacs in X–XI. Sv paired in XI–XII. Prostates in XVIII, extending between XVII–XX; prostatic ducts tightly folded twice. Genital marking glands paired in XVII and XVIII corresponding to external genital papillae, each consisting of a stalk with terminal multi-lobed glandular part.

Ovaries in XIII. Sc one pair in VIII; ampulla oval to kidney-shaped, with stout duct shorter than ampulla. Diverticulum with oval bulb terminal, stalk attached to duct near body wall. Genital markings stalked, corresponding to external genital papillae; each gland small consisting of a stalk with terminal multi-lobed glandular part.

Table 5. Holotype and Paratype dimension and other morphological characteristics of *Amyntas srinan* Panha & Bantaowong, sp. n.

Types	Characters	Body length (mm)	Number of segments	Location of genital markings		First dorsal pore	Number of setae		Between male pore	Prostate glands	Intestinal caeca
				preclitellum	postclitellum		VII	XX			
Holotype		47	77	VII, VIII	XVII, XVIII	5/6	36	42	4	XVII-XX	XXVII-XXIII
CUMZ 3210											
Paratype											
CUMZ 3211											
1		35	75	VII, VIII	XVII, XVIII	5/6	40	42	6	XVII-XX	XXVII-XXV
2		44	76	VII, VIII	XVII, XVIII	5/6	36	42	5	XVII-XX	XXVII-XXIV
3		39	65	VII, VIII	XVII, XVIII	5/6	37	46	4	XVIII-XX	XXVII-XXIV
4		44	70	VII, VIII	XVII, XVIII	5/6	36	49	5	XVII-XIX	XXVII-XXIV
5		47	78	VII, VIII	XVII, XVIII	5/6	38	45	4	XVII-XX	XXVII-XXIV
6		37	68	VII, VIII	XVII, XVIII	5/6	40	44	4	XVII-XX	XXVII-XXV
7		38	77	VII, VIII	XVII, XVIII	4/5	43	48	5	XVII-XXI	XXVII-XXIV
8		37	52	VII, VIII	XVII, XVIII	4/5	38	42	4	XVII-XXI	XXVII-XXV
9		35	57	VII, VIII	XVII, XVIII	4/5	41	44	4	XVII-XX	XXVII-XXIV
10		38	78	VII, VIII	XVII, XVIII	5/6	36	40	4	XVII-XX	XXVII-XXIV
11		42	77	VII, VIII	XVII, XVIII	4/5	42	47	4	XVII-XXI	XXVII-XXIII
12		45	77	VII, VIII	XVII, XVIII	5/6	39	45	5	XVII-XX	XXVII-XXIV
13		40	77	VII, VIII	XVII, XVIII	5/6	40	48	4	XVII-XIX	XXVII-XXV
14		39	77	VII, VIII	XVII, XVIII	5/6	39	47	4	XVII-XX	XXVII-XXIV
15		43	77	VII, VIII	XVII, XVIII	5/6	40	44	4	XVII-XX	XXVII-XXIII
16		40	75	VII, VIII	XVII, XVIII	5/6	41	49	4	XVII-XX	XXVII-XXIV
17		37	75	VII, VIII	XVII, XVIII	4/5	36	46	4	XVII-XIX	XXVII-XXIV
18		36	60	VII, VIII	XVII, XVIII	5/6	40	47	5	XVII-XX	XXVII-XXIV
19		39	75	VII, VIII	XVII, XVIII	5/6	37	44	4	XVII-XX	XXVII-XXIII

Characters Types	Body length (mm)	Number of segments	Location of genital markings		First dorsal pore	Number of setae		Between male pore	Prostate glands	Intestinal caeca
			preclitellum	postclitellum		VII	XX			
20	47	78	VII, VIII	XVII, XVIII	4/5	36	42	4	XVII–XX	XXVII–XXV
21	42	71	VII, VIII	XVII, XVIII	5/6	40	46	4	XVII–XX	XXVII–XXIV
22	35	56	VII, VIII	XVII, XVIII	4/5	41	43	4	XVII–XIX	XXVII–XXIV
23	36	69	VII, VIII	XVII, XVIII	5/6	36	45	4	XVII–XX	XXVII–XXV
24	42	73	VII, VIII	XVII, XVIII	5/6	36	46	4	XVII–XX	XXVII–XXIV
25	44	76	VII, VIII	XVII, XVIII	4/5	39	47	6	XVI–XX	XXVII–XXV
26	35	69	VII, VIII	XVII, XVIII	5/6	36	44	4	XVII–XIX	XXVII–XXIII
27	38	75	VII, VIII	XVII, XVIII	5/6	37	45	4	XVII–XX	XXVII–XXIV
28	35	78	VII, VIII	XVII, XVIII	5/6	39	44	4	XVII–XIX	XXVII–XXV

All the key morphological characters of the holotype and paratype specimens are given in Table 5.

Variation: The holotype measures 47 mm body length with 77 segments and the first dorsal pore located at 5/6; the twenty eight paratypes range in size between 35–47 mm (39.75 ± 4.27 mm) body length with 52–78 segments, and first dorsal pore at 4/5 (8 samples) or 5/6 (20 samples) (Table 5).

Type locality: Srinan National Park, Nan province, Thailand, $18^{\circ}22'11.1''\text{N}$, $100^{\circ}50'23.2''\text{E}$, 607 meters elevation (30th September 2010).

Etymology: This species was named after the type locality Srinan National Park.

Type material: The holotype (CUMZ 3210) and 25 paratypes (CUMZ 3211) are deposited in Chulalongkorn University, Museum of Zoology. Another five paratypes will be deposited in the Biozentrum Grindel und Zoologisches Museum, Hamburg, Germany (UHH), and four paratypes in the Natural History Museum, London (NHM).

Habitat: Found in the top soil at about 10 cm depth, the soil surface covered with leaf litters of deciduous forest. The soil was carefully dug close to the castes.

Diagnosis: *Amyntas srinan* sp. n. is the smallest *Amyntas* ever collected in Thailand. Male pores on distinct round porophores, genital markings paired near mid ventral of VII, VIII, XVII and XVIII; each with genital marking glands. Each spermathecae consists of a kidney-shaped ampulla and an oval shaped diverticulum. Testes sacs ventrally joined, intestinal origin XV, intestinal caeca simple, first dorsal pores at 4/5 or 5/6.

Remarks: *Amyntas srinan* sp. n., along with *A. exiguus exiguus* and *A. tontong* sp. n., is one of if not the smallest *Amyntas* recorded so far. It has external characteristics which are easily seen in both newly collected and preserved materials. Compared with the two other closely related species from Laos, *A. chandyi* and *A. namphouinensis*, which belong in the same *zebrus*-group, *A. chandyi* is very similar in appearance to *A. srinan* sp. n. However, they clearly differ in certain specific details of their significant characters, such as the distance between the mp which in *A. srinan* sp. n. is 1.5 mm for holotype and ranged from 1.5–2.0 mm (1.41 ± 4.27 mm), while in *A. chandyi* this ranged from 1.5–2.4 mm. The distance between the male pores as a fraction of the estimated circumference of the 18th segment is 0.24–0.30 in *A. srinan* sp. n., and 0.14–0.32 in *A. chandyi*. This is not convincing as a diagnostic difference, because there is significant overlap with the highly variable *A. chandyi*. In addition, although genital markings are clearly observed in both *A. chandyi* and *A. srinan* sp. n. on the sc and mp areas, *A. srinan* sp. n. has a much larger number and different arrangement of such markings. The distance between pairs of sp is quite similar, being 1.5–2.0 mm (1.34 ± 2.31 mm) in *A. srinan* sp. n. and 1.2–1.5 mm in *A. chandyi*.

Discussion

The genus *Amyntas* is widely distributed in the Asian continent, where it is one of the dominant genera. In Thailand it occurs in various types of lowland forest habi-

tats, dry evergreen, moist evergreen, deciduous and limestone forests, encompassing diverse soil pH values, from acidic to alkali soils (Chantaravisoot, 2007) and from clay to muddy sand substrates (Kosavitikul, 2005; Somniyam, 2008; Blakemore et al., 2007). The current four new species described here were all found in one area (Nan province) but the four habitat types were quite diverse all the same. *Amyntas phatubensis* sp. n. was found in a limestone area with a mild alkali substrate (pH 7.5–8) of a clay loam structure, whilst the other three species were found in harder sandy clay substrates. The four new species are broadly similar (and so potentially related) to the two species described from Laos, *A. chandyi* and *A. namphouinensis*, but differ in both the external and internal morphological characteristics. The geographic structures of Luang Prabang Mountain and Phi Pan Nam Mountain ranges are important barriers for species from both the Thai (Nan province) and Laos side (Xayabouli province) and may have played an important part in their speciation. In addition, the Laos species live at a higher altitude than the current new described species from Thailand, and such selective adaptations may facilitate their morphological discrimination.

The four new species range in size, with respect to other *Amyntas* members, from moderate to very small, of which *A. phatubensis* sp. n. is the longest. The other three species are almost the same size and close to the two Laotian species, as shown in Table 6. However, the spermathecae (sc) and genital marking locations of the four new species are clearly different from the two closely related Laos species. The four new *Amyntas* species described here belong to the *zebrus*-group, as defined by Sims and Easton (1972), in which the spermathecal pores are located on segment 7/8. The size of these four species, relative to other *Amyntas* species, varied from small to medium, ranging from 35 to 148 mm in body length and having from 52 to 114 segments. The first dorsal pore in three of the four species described here, and most of the samples of the fourth species (*A. srinan* sp. n.), is located on intersegmental furrow 5/6, but with some samples of *A. srinan* sp. n. showing the first dorsal pore at 4/5.

Amyntas phatubensis sp. n. is the only species that lives in limestone habitats in leaf litter and also in shallow mild alkali topsoil. The soil humidity can be quite low and is of a clay loam structure. The other three species are smaller in size and were found in almost harder, muddy sandy clay substrates. *Amyntas tontong* sp. n. lives in deeper soil of a high humidity around waterfalls. *Amyntas borealis* sp. n. and *A. srinan* sp. n. are found in deciduous forests, which have mostly been modified as agricultural fields. The soil is drier and harder. The genital marking glands of *A. phatubensis* sp. n. and *A. srinan* sp. n. are distinct from other two species (Table 6 and Figs 2–5), whilst *A. tontong* sp. n. show two postclitellar genital markings that are absent in *A. borealis* (Figs 3 and 4) The diagnostic differences are shown in the dichotomous key to the sixteen Thai and two Laotian *Amyntas* species, below.

The *zebrus*-group is composed of eleven nominal species: *Metaphire hilgendorfi* (Michaelsen, 1892), *A. palmosus* (Chen, 1946), *A. magnipapillatus* (Qui and Wang, 1992), *A. zebrus* (Benham, 1896), *A. culminus* Michaelsen, 1899, *A. principalis* (Michaelsen, 1932), *A. xuongmontis* (Thai & Samphon, 1990), *A. fasciculus* (Qui,

Table 6. Morphological characteristics for between these four new species and two know species from Laos

Characters	<i>A. phatubensis</i> sp. n.	<i>A. tontong</i> sp. n.	<i>A. borealis</i> sp. n.	<i>A. srinan</i> sp. n.	<i>A. namphouinensis</i>	<i>A. chandyi</i>
Body length (mm)	80–148	39–53	42–54	35–47	63–92	29–58
Number of segments	85–112	71–80	78–89	56–77	92–94	48–52
First dorsal pore	5/6	5/6	5/6	4/5, 5/6	4/5, 5/6, 6/7	5/6
Setae number						
VII	51–64	41–46	39–54	36–45	52–61	44–54
XX	58–68	52–55	40–52	42–49	53–58	44–57
between male pores	9–15	0	0	4–6	0–7	0–7
Precitellar genital markings						
VII	2	0	0	2	0	1–2
VIII	1–7	0	0	2	0	1–2
IX	0–1	0	0	0	0	0
Postclitellar genital markings						
XVII	0–2	0	0	2	2	1
XVIII	6–12	2	0	2	0	3
XIX	0–1	0	0	0	4	1
XX	0–1	0	0	0	0	1
Prostate glands	XVII–XX	XVII–XX	XVII–XX	XVII–XX	XVII–XIX	XVI–XXI
Genital marking glands	sessile at VII, VIII	Absent	absent	stalked	sessile at XVII–XIX	absent
Intestinal caeca	simple, XXVII–XXIII	simple, XXVII–XXV	simple, XXVII–XXV	simple, XXVII–XXII	simple, XXVII–XXIV	simple, XXVII–XXIV




Figure 6. External and internal morphology of holotype (BDNUL 0001) of *Amynthus namphouinensis* Hong, 2008 **A** External ventral view, **B** internal dorsal view and **C** spermatheca, and black arrow indicates the connection of the spermatheca and spermathecal pore.

Wang & Wang, 1993), *A. heaneyi* James, 2004, *A. namphouinensis* Hong, 2008 and *A. chandyi* Hong, 2008. Within the *zebrus*-group, the first three species show manicate intestinal caeca, while the current newly described four species have simple finger-shaped intestinal caeca. The three latter nominal species are longer in body length (200–300 mm) compared with the size of these four new species which ranged from 35–148 mm. *Amynthus heaneyi* can be distinguished by its proandric character (James, 2004), while the four new described species are holandric. *Amynthus fasciculus* has coiled and kinked spermathecae, whereas *A. phatubensis* sp. n. has large ovate ampulla, *A. tontong* sp. n. has thumb shaped ampulla, *A. borealis* sp. n. has sac-shape ampulla, and *A. srinan* sp. n. has oval to kidney-shaped ampulla. *Amynthus xuongmontis* clearly differs from these four new species in the genital marking located on XVIII, whereas located on VII, VIII, XVII, XVIII in *A. phatubensis* sp. n., located between 18/19 in *A. tontong* sp. n., absent in *A. borealis* sp. n. and located on VII, VIII, XVII, XVIII in *A. srinan* sp. n.




Figure 7. External and internal morphology of holotype (BDNUL 0002) of *Amynthes chandyi* Hong, 2008 **A** External ventral view, **B** internal dorsal view and **C** spermatheca, and black arrow indicates the connection of the spermatheca and spermathecal pore.

Key to Thai and two Laos species of *Amynthes*

- 1
- First spermathecal pores at 5/6.....2
-
- First spermathecal pores after 5/6.....12
- 2
- Two pairs of spermathecal pores.....*A. morrisi*
-
- More than two pairs of spermathecal pores3
- 3
- Three pairs of spermathecal pores.....4
-
- More than three pairs of spermathecal pores6
- 4
- Genital markings absent*A. defecta*
-
- Genital markings present5
- 5
- Genital markings clustered on XVIII*A. gracilis*
-
- Genital markings transverse rows on XVII, XVIII, XIX*A. papulosus*
- 6
- Genital markings absent7
-
- Genital markings present8
- 7
- Body length 1 meter or more*A. mekongianus*

–	Body length less than 300 mm.....	<i>A. alexandri</i>
8	Genital marking glands absent.....	9
–	Genital marking glands present.....	10
9	Genital markings located on 17/18, 18/19.....	<i>A. exiguus austrinus</i>
–	Genital markings located on VII, VIII, XIX, XX	<i>A. exiguus exiguus</i>
10	Intestinal caeca, simple	11
–	Intestinal caeca, manicate.....	<i>A. manicatus decorosus</i>
11	Genital markings, paired at 18/19, 19/20, 20/21	<i>A. longicauliculatus</i>
–	Genital markings, three trios at 18/19, 19/20, 20/21	<i>A. comptus</i>
12	First spermathecal pores at 6/7.....	13
–	First spermathecal pores after 6/7.....	14
13	Genital markings located on 17/18, 18/19.....	<i>A. fucosus</i>
–	Genital marking located on XVIII	<i>A. siam</i>
14	Body length more than 200 mm.....	<i>A. hupbonensis</i>
–	Body length less than 200 mm.....	15
15	Genital markings absent	<i>A. borealis sp. n.</i>
–	Genital markings present	16
16	Preclitellar genital markings absent	17
–	Preclitellar genital markings present.....	18
17	Genital marking glands absent.....	<i>A. tontong sp. n.</i>
–	Genital marking glands present.....	<i>A. namphouinensis</i>
18	Genital marking glands absent.....	<i>A. chandyi</i>
–	Genital marking glands, present.....	19
19	Genital marking glands, sessile.....	<i>A. phatubensis sp. n.</i>
–	Genital marking glands, stalked	<i>A. srinan sp. n.</i>

Acknowledgements

This project was the Higher Education Research Promotion and National Research University Project of Thailand, Office of the Higher Education Commission FW646A (2011–2013) to SP, and from Science for Locale Project, Faculty of Science, Chulalongkorn University. We are grateful to Khamla Inkhavilay for kindly permitting us to study the type specimens and relevant reference material of National University of Laos at Vientiane. Thanks also to Thita Krutchuen for excellent drawings, and to all members of the Animal Systematics Research Unit, Chulalongkorn University for assistance in collecting material.

References

- Beddard FE (1892) On some species of the genus *Perichaeta* (sensu strictu). Proceedings of the Zoological Society of London 1892: 153–172.
- Beddard FE (1900) On a new species of earthworm from India belonging to the genus *Amyntas*. Proceedings of the Zoological Society of London 1900: 998–1002.
- Blakemore RJ (2000) ‘Tasmanian Earthworms.’ CD-ROM Monograph with review of world families. VermEcology, Kippax, Canberra, December 2000, 800 pp.
- Blakemore RJ (2006a) Introductory key to the Revised Families of Earthworms of the world. In: Kaneko N, Ito MT (Eds) A Series of Searchable Texts on Earthworm Biodiversity, Ecology and Systematics from Various Regions of the World. CD-ROM publication by Soil Ecology Research Group, Graduate School of Environment & Information Sciences, Yokohama National University, Yokohama, Japan, 14 pp. [accessed 3 April 2011]
- Blakemore RJ (2006b) Checklist of Thailand taxa updated from Gates’ (1939) “Thai Earthworms”. In: Ito MT, Kaneko N (Eds) A Series of Searchable Texts on Earthworm Biodiversity, Ecology and Systematics from Various Regions of the World. CD-ROM publication by Soil Ecology Research Group, Graduate School of Environment & Information Sciences, Yokohama National University, Yokohama, Japan, 6 pp. [accessed 21 December 2010]
- Blakemore RJ, Csuzdi C, Ito MT, Kaneko N, Paoletti MG, Spiridonov SE, Uchida T, Van Praagh BD (2007) *Megascolex* (*Promegascolex*) *mekongianus* Cognetti, 1922: its extent, ecology and allocation to *Amyntas* (Oligochaeta: Megascolecidae). Opuscula Zoologica 36: 19–30.
- Blakemore RJ (2008) Cosmopolitan Earthworms – an Eco-Taxonomic Guide to the Peregrine Species of the World (3rd Edition). Verm Ecology, Yokohama, Japan, 757 pp.
- Blakemore RJ, Kupriyanova EK, Grygier MJ (2010) Neotypification of *Drawida hattamimizu* Hatai, 1930 (Annelida, Oligochaeta, Megadrili, Moniligastridae) as a model linking mtDNA (COI) sequences to an earthworm type, with a response to the ‘Can of Worms’ theory of cryptic species. Zookeys 41: 1–29.
- Blakemore RJ (2011) Description of a new *Amyntas* earthworm (Megascolecidae *sensu stricto*) from Thailand. Bulletin of the National Science Museum Series A 37: 9–13.
- Chang C–H, Lin Y–H, Chen I–H, Chuang S–C, Chen J–H (2007) Taxonomic re-evaluation of the Taiwanese montane earthworm *Amyntas wulinensis* Tsai, Shen & Tsai, 2001 (Oligochaeta: Megascolecidae): polytypic species or species complex?. Organism Diversity & Evolution 7: 231–240.
- Chantaravisoot N (2007) Species Diversity of Terrestrial Earthworms Family Megascolecidae in Thailand. Senior Project Report, Department of Biology, Faculty of Science, Chulalongkorn University, 36 pp. [with English Abstract]
- Folmer O, Back M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome *c* oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology and Biotechnology 3:294–299.
- Gates GE (1930) The earthworms of Burma I. Records of the Indian Museum 32: 257–356.
- Gates GE (1931) The earthworms of Burma II. Records of the Indian Museum 33: 327–442.
- Gates GE (1932) The earthworms of Burma III. Records of the Indian Museum 34: 357–549.

- Gates GE (1933) The earthworms of Burma IV. Records of the Indian Museum 35: 413–606.
- Gates GE (1972) Burmese earthworms, an introduction to the systematics and biology of megadrile oligochaetes with special reference to the Southeast Asia. Transactions of the American Philosophical Society 62: 1–326.
- Hajibabaei M, deWaard JR, Ivanova NV, et al. (2005) Critical factors for assembling a high Volume of DNA barcodes. Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences 360: 1959–1967.
- Hebert PDN, Cywinska A, Ball SL, deWaard JR (2003a) Biological identifications through DNA barcodes. Proceedings of the Royal Society of London. Series B: Biological science 270: 313–321.
- Hebert PDN, Ratnasingham S, deWaard JR (2003b) Barcoding animal life: cytochrome *c* oxidase subunit 1 divergences among closely related species. Proceedings of the Royal Society of London. Series B: Biological science 270(Suppl.): S96–S99.
- Hong Y (2008) Two new bithecal earthworms of the genus *Amyntas* (Oligochaeta: Megascollecidae) from Laos. Zootaxa 1914: 57–61.
- Ivanova NV, Jeremy R, deWaard, Hebert PDN (2006) Technical note: An inexpensive, automation-friendly protocol for recovering high-quality DNA. Molecular Ecology Notes 6: 998–1002
- James SW (2004) New species of *Amyntas*, *Pheretima* and *Pleionogaster* (Clitellata: Megascollecidae) of the Mt. Kitanglad Range, Mindanao Island, Philippines. The Raffles Bulletin of Zoology 52: 289–313.
- James SW, Porco D, Decaëns T, Richard B, Rougerie R, et al. (2010) DNA Barcoding Reveals Cryptic Diversity in *Lumbricus terrestris* L., 1758 (Clitellata): Resurrection of *L. herculeus* (Savigny, 1826). PLoS ONE 5(12): e15629. doi:10.1371/journal.pone.0015629
- Kimura M (1980) A Simple Method for Estimating Evolutionary Rates of Base Substitutions Through Comparative Studies of Nucleotide Sequences. Journal of Molecular Evolution 16: 111–120
- Kosavittikul P (2005) Species Diversity of Terrestrial Earthworms in Khao Yai National Park. Ph.D Thesis in Environmental Biology, School of Biology, Institute of Science, Suranaree University of Technology, 193 pp.
- Pérez-Losada M, Eiroa J, Mato S, Domínguez J (2005) Phylogenetic species delimitation of the earthworm *Eisenia fetida* (Savigny, 1826) and *Eisenia andrei* (Bouché, 1972) (Oligochaeta, Lumbricidae) based on mitochondrial and nuclear DNA genes, Pedobiologia 49: 317–324.
- Ratnasingham S, Hebert PND (2007) Barcoding Bold: The Barcode of Life Data System (www.barcodinglife.org). Molecular Ecology Notes 7: 355–364.
- Rosa D (1891) Die exotoschen Terricolen des k.k. naturhistorischen Hofmuseums. Annalen des k.k. Naturhistorischen Hofmuseums, Wein 6: 379–406.
- Rosa D (1896) I Lombrichi raccolti a Sumatra dal dott elio modigliani. Annali del Museo Civico di Storia Naturale, 2nd Serie 16: 502–532.
- Saitou N, Nei M (1987) The Neighbor-joining Method: A New Method for Reconstructing Phylogenetic Trees. Molecular Biology and Evolution 4: 406–425.
- Shen H-P, Yeo DCJ (2005) Terrestrial earthworms (Oligochaeta) from Singapore. The Raffles Bulletin of Zoology 53: 13–35.

- Sims RW, Easton EG (1972) A numerical revision of the earthworm genus *Pheretima* (Megascolecidae: Oligochaeta) with the recognition of new genera and an appendix on the earthworms collected by the Royal Society North Borneo Expedition. *Biological Journal of the Linnean Society* 4: 169–268.
- Somniyam P (2008) The Population Dynamics and Distribution of Terrestrial Earthworms at Sakaerat Environmental Research Station and Adjacent Areas, Nakornratchasima Province. Ph.D Thesis in Environmental Biology, School of Biology, Institute of Science, Suranaree University of Technology, 232 pp.
- Stephenson J (1931) Oligochaeta from Burma, Kenya and other parts of the world. *Proceedings of the Zoological Society of London* 1931: 33–92.
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* 24: 1596–1599.

Appendix I.

DNA barcode sequences for *A. phatubensis* sp. n. and *A. tontong* sp. n. Positions with variable base are indicated by the appropriate ambiguity code: Y= C or T, R= A or G, K= G or T, M= A or C.

The primer sets used, LCO1490 and HCO2198, amplify a 658 bp fragment of the COI gene in a wide range of invertebrate taxa (Folmer et al 1994):

LCO1490: 5'-GGTCAACAAATCATAAAGATATTGG-3'

HCO2198: 5'-TAACTTCAGGGTGACCAAAAAATCA-3'

Consensus of 8 sequences of *Amyntas phatubensis* sp. n. **Paratype** CUMZ 3212, GenBank Accession No. HM901031-HM901038.

AACCCTATATTTCATTTT TAGGTATTTGAGCCGGTATGATTGGATCTG-
GAATAAGCCTACTYATCCGAATTGARTTGAGCCAACCTGGATCCTTC-
CTAGGCAGAGATCAGCTATAACAATACCATTTGTTACAGCTCATGCATTCT-
TAATAATYTTCTTTTTTAGTTATACCCGTATTTATTGGGGGGTTTTG-
GAAACTGATTAYKACCACTTATACTTGGRGCGCCAGATATRGCCCTTTC-
CYCGACTAAACAACATAAGATTCTGATTGCTTCCTCCRTCRCCTTATTC-
TATTAGTAAGCTCTGCGGGCCGTRGAAAAGGGGGGCGYGGCACTGGAT-
GAACTGTTTACCCMCCCCCTAGCTAGAAATGTRGCCCACGCAGGGC-
CTTCAGTAGATTTAGCTATTTTCTCACTTCATTTAGCAGGAGCTTCATC-
TATTTTAGGGGGCAATTAATTTTCAATTACAACCTGTAATTAATATGCGAT-
GATCCGGACTACGTCTAGAGCGTATCCCACCTATTTGTATGGGC-
CGTRGTTATCACAGTAGTACTATTATTACTTTCCCTRCCTGTATT-
AGCAGGGGGCTATCACTATACTACTAACTGAYCGTAATCTAAACACAT-
CATTTTTTGATCCTGCTGGGGGCGGGCGACCCCATTCATATCAACAC-
CTA

Consensus of 3 sequences of *Amyntas tontong* sp. n. **Paratype** CUMZ 3207, GenBank Accession No. HQ562073-HQ562076.

AACCCTATACTTCATTTT TAGGAATTTGAGCTGGAATAATTGGAGCAG-
GAATAAGACTCCTTATTCGAATTGAGYTAAGACAGCCCCGGATCATTC-
CTAGGAAGYGATCAACTATAACAATACCATTGTTACAGCCCATGCATTCT-
TAATAATTTTCTTCCTRGTAATGCCAGTATTTATTGGGGGGCTTTG-
GAAATTGATTACTTCCACTAATGTTGGGGGGCCCCCTGACATAGCTTTC-
CCACGACTAAATAATATGAGATTTTGTACTACTTCCACCATCATTAATC-
CTTTTAGTTAGATCCGCRGCCGTTGAAAAAGGTGCGGGGGACAGGAT-
GAACTGTATACCCACCACTAGCAAGGAACATTGCCCCATGCTGGCC-
CATCTGTAGATTTAGCAATTTTCTCACTACACTTGGCGGGGGGCATCCT-
CAATCCTGGGGGGCTATTA ACTTCATTACCACAGTAATTAATATGCGAT-
GATCTGGGCTGCGCTTAGAACGAATCCCCCTATTTGTATGAGCTGTAG-
TAATTACAGTAGTACTCCTACTACTATCTTTGCCCGTGCTGGCGGGGAGC-
CATTACAATACTCTTAACAGATCGAAATCTTAATACATCATTCCTTCGACC-
CTGCTGGCGGGGGGCGACCCTATTCTATACCAGCACCTG